

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**


333/NF/03

We Claim:


- 1 Pyrrolo[2,1-c][1,4]benzodiazepine hybrid of the formula given below wherein R is H or OH and n is 2-3




$n = 2-3$
 $R = H, OH$




- 5 2 Pyrrolobenzodiazepine hybrid as claimed in claim 1 of the structure



- 3 Pyrrolobenzodiazepine hybrid as claimed in claim 1 of the structure




- 4 Pyrrolobenzodiazepine hybrid as claimed in claim 1 of the structure




15


- 5 Pyrrolobenzodiazepine hybrid as claimed in claim 1 of the structure




- 6 Pyrrolobenzodiazepine hybrid as claimed in claim 1 of the structure




- 5 7 Pyrrolobenzodiazepine hybrid as claimed in claim 1 of the structure




- 8 Pyrrolobenzodiazepine hybrid as claimed in claim 1 of the structure




- 9 Pyrrolobenzodiazepine hybrid as claimed in claim 1 of the structure



- 10 A process for the preparation of a compound of the formula wherein R is H or OH and n is 2-3




the process comprising reacting reacting an acridone or an acridine acid with (2*S*)-N-[4-(*n*'-aminoalkoxy)-5-methoxy-2-nitrobenzoyl]-pyrrolidine-2-carboxaldehyde diethyl thioacetal of formula I




Formula I

in the presence of EDCI and HOBt in organic solvent for a period of 24 h to obtain (2*S*)-N-{4-[*n*'-(4''-acrido-nylcarboxamido)-alkyl]-oxy-5-methoxy-2-nitrobenzoyl} pyrrolidine-2-carboxaldehyde diethyl thioacetal II / (2*S*)-N-{4-[*n*'-(4''-acridinylcarboxamido)-alkyl]-oxy-5-methoxy-2-nitrobenzoyl}pyrrolidine-2-carboxaldehyde diethyl thioacetal of formula V where '*n*' is 2-3,




Formula II




Formula V

isolating the compound of formula II/formula V and then reducing the compounds of formula II/formula V with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ in presence of an organic solvent up to a reflux temperature, isolating the (2*S*)-N-{4-[n'-(4''-acridonylcarboxamido)-alkyl]-oxy-5-methoxy-2-aminobenzoyl}pyrrolidine-2-carboxaldehydediethylthioacetal of formula III/(2*S*)-N-{4-[n'-(4''-acridinylcarbox-amido)-alkyl]-oxy-5-methoxy-2-aminobenzoyl}pyrrolidine-2-carboxaldehyde diethyl thioacetal of formula VI where n is 2-3,



Formula III



Formula VI

reacting compound of formula III/formula VI with a deprotecting agent to obtain the desired pyrrolo[2,1-*c*][1,4]benzodiazepine hybrid.

A process as claimed in claim 10 wherein the organic solvent used for the reaction of the acridone/acridine acid with compound of formula I comprises dimethyl furan.

A process as claimed in claim 10 wherein the compound of formula II/formula V is isolated by washing with saturated NaHCO_3 , brine, drying and evaporation of the solvent.


A process as claimed in claim 10 wherein the organic solvent used during the reduction of compound of formula II/formula V comprises methanol.

A process as claimed in claim 10 wherein the compound of formula III/formula V is isolated by adjusting the pH of the reaction mixture to about pH 8 with a saturated


NaHCO₃ solution, diluting with ethyl acetate, filtering through celite and extracted an organic phase and drying the organic phase over Na₂SO₄.

15 A process as claimed in claim 10 wherein the deprotecting agent used for obtaining the compound of formula IV/formula VII comprises HgCl₂ and CaCO₃ in MeCN-water (4:1).

16 A pharmaceutical composition comprising a pharmaceutically effective amount of a compound of the formula given below wherein R is H or OH and n is 2-3 and a pharmaceutically acceptable additive.



17 A method for the treatment of cancer in a subject suffering from the same comprising administering a pharmaceutically effective amount of a compound of the formula



wherein R is H or OH and n is 2-3.

18. A method as claimed in claim 17 wherein the patient is a mammal.
19. A method as claimed in claim 17 wherein the mammal is a human being.
20. A method as claimed in claim 17 wherein the cancer is selected from the group consisting of leukemia, non-small cell, lung, colon, CNS, melanoma, ovarian, renal, prostate and breast.

21. Use of a compound of formula given below for the treatment of cancer selected from the group consisting of leukemia, non-small cell, lung, colon, CNS, melanoma, ovarian, renal, prostate and breast in a subject suffering from the same.

